

9/23/08

(1)

Digression on quaternions. Quaternions don't have anything to do with homotopy, but they are related to  $SU(2)$ , so we'll say something about them now. Recall any  $u \in SU(2)$  can be written

$$u = x_0 \mathbf{1} - i \vec{x} \cdot \vec{\sigma}$$

where  $(x_0, x_1, x_2, x_3)$  are the Cayley-Klein parameters and  $\sum_{i=0}^3 x_i^2 = 1$ .

Thus every element of  $SU(2)$  corresponds to a unit vector in  $\mathbb{R}^4$  and vice versa, and  $SU(2) \cong S^3 \subset \mathbb{R}^4$ .

We get the quaternions if we drop the constraint on the 4  $x$ 's and let them run all over  $\mathbb{R}^4$ . Thus if  $q \in \mathbb{H}$  (the set of quaternions), then

$$q = x_0 \mathbf{1} - i \vec{x} \cdot \vec{\sigma}$$

and  $\mathbb{H} \cong \mathbb{R}^4$ , with a multiplication rule given by the expression above and the algebra of Pauli matrices. Sometimes this is written

$$q = x_0 \mathbf{1} + x_1 \hat{i} + x_2 \hat{j} + x_3 \hat{k},$$

where  $\hat{i}^2 = \hat{j}^2 = \hat{k}^2 = -1$ . Here in a sense we have a representation of the quaternions by means of  $2 \times 2$  matrices.

Obviously any vector in  $\mathbb{R}^4$  can be written as a magnitude  $r$  times a unit vector, so

$$q = r u, \quad \begin{matrix} 0 \leq r \\ u \in SU(2) \end{matrix}$$


The elements of  $SU(2)$  are the unit quaternions. This is like the complex numbers, every  $z \in \mathbb{C}$  can be written  $z = r e^{i\phi}$  where  $0 \leq r$  and  $e^{i\phi} \in U(1)$ . Multiplication of quaternions is noncommutative since  $SU(2)$  is Abelian.

9/23/08

(2)

Return to  $SU(2)$  as a cover of  $SO(3)$ . Let's divorce ourselves from the eqns. of evolution for  $R(t)$  and  $U(t)$  and look at topological aspects.

Not a very realistic representation of the relation betw.  $SU(2)$  and  $SO(3)$



(contin.)

Let  $R(t)$  be a path in  $SO(3)$  starting at  $R(0) = I$ . We wish to find a path  $U(t)$  in  $SU(2)$  such that  $p: U(t) \mapsto R(t)$ . Since  $p^{-1}$  is double-valued, there is no unique answer. But if we require that  $U(0) = 1$  and that  $U(t)$  be continuous, then the answer is unique. First, ~~unless~~  $U(0) = +1$  picks one of the two preimages at  $t=0$ . Then, for each small step we make in  $R(t)$ , only one of the two possible preimages is possible if we demand  $U(t)$  be continuous, because one of the preimages is close to the part of  $U(t)$  developed already and the other is far away. Note that  $U$  and  $-U$  are on opposite sides of  $S^3$  and never come close together. The path  $U(t)$  created in this way is the lift of  $R(t)$ . It is like the lifts we will see later in fiber bundle theory, except the law of "parallel transport" is determined by continuity.

$$R(0) =$$

Now let  $R(t)$  be a loop, i.e.  $R(T) = I$ . This loop belongs either to the trivial or nontrivial class of  $\pi_1(SO(3)) = \mathbb{Z}_2$ . We now show that  $R(t)$  is trivial iff  $U(T) = +1$ , where  $U(t)$  is

9/23/08 ③

the lift of  $R(t)$ . This tells us the state of the neutron after the classical rotation has returned to the identity.

If  $R(t)$  is contractable (trivial), we wish to show that  $U(T) = +1$ . Suppose not, i.e. suppose  $U(T) = -1$ . Then the lift of the loop  $R(t)$  is an open path  $U(t)$  in  $SU(2)$  that ends at  $U(T) = -1$ . Now as we contract  $R(t)$  down to the constant loop  $c: I \rightarrow SO(3): t \mapsto Id$ ,  
 the lift of the contracting <sup>loop</sup>~~curve~~ is a path  $U(t)$  that must end  $\hookrightarrow [0, T]$   
 on either  $+1$  or  $-1$ , since  $R(T) = I$ . At the beginning of the  
 contraction,  $U(T) = -1$  (we are supposing), while at the end ~~it~~ we  
 have the lift of the constant loop in  $SO(3)$  which is the constant  
 loop:  $[0, T] \rightarrow SU(2): t \mapsto 1$  in  $SU(2)$ . But the endpoint cannot  
 change continuously from  $-1$  to  $+1$ , while the lifting process  
 is continuous. A continuous function that takes values in a discrete  
 set must be constant. So the assumption  $U(T) = -1$  must be wrong,  
 and we must have  $U(T) = +1$ .

Conversely, suppose we have a loop  $R(t)$  on  $SO(3)$  that lifts into a loop  $U(t)$  on  $SU(2)$ , i.e. a path that ends in  $U(T) = +1$ . Since  $SU(2) \cong S^3$  is simply connected, this loop is contractable. As it contracts, its projection onto  $SO(3)$  contracts continuously to the constant loop. Therefore  $R(t)$  belonged to the trivial class.

9/23/08

(4)

Now we give the official definition of a covering space.


Def. Let  $M, \bar{M}$  be connected topological spaces with a map  $p: \bar{M} \rightarrow M$  such that

- (1)  $p$  is surjective
- (2) for each  $x \in M \exists$  a connected open neighborhood  $V \subset M$  containing  $x$  such that  $p^{-1}(V)$  is a disjoint union of open sets  $\{V_\alpha\}$  in  $\bar{M}$ , each mapped homeomorphically onto  $V$  by  $p$ ,  $p(V_\alpha) = V, \forall \alpha$ .


$(M = \text{"original" or "base" space})$   
 $(\bar{M} = \text{"covering" space})$

Then  $\bar{M}$  is ~~the~~ a covering space of  $M$ . If  $\bar{M}$  is simply connected, then it is the universal covering space.

Remarks. we require  $M$  to be connected, because otherwise we might as well talk about the cover of each component (piece) of  $M$  separately:



And we require  $\bar{M}$  to be connected, because otherwise  $M$  can be thought of as having more than one cover, each of which can be treated separately:




9/23/08

(5)

Look at requirement (2), regarding open sets, in the case of  $SU(2)$  covering  $SO(3)$ . Represent  $SO(3)$  as the northern hemisphere of  $S^3$ , let  $R \in SO(3)$ , and let  $V$  be a neighborhood of it:



Then look at  $p^{-1}(V)$  on  $SU(2)$ , represented as the whole of  $S^3$ :



$$p^{-1}(V) = V_1 \cup V_2$$

$$V_1 \cap V_2 = \emptyset$$

It means that in the neighborhood of each pre-image of  $p^{-1}(R)$ ,  $SU(2)$  "looks like"  $SO(3)$ , topologically speaking. The fact that  $V_1 \cap V_2 = \emptyset$  means that the preimages  $\overset{(u, -u)}{\curvearrowright}$  are well separated from each other, and remain so as  $R$  moves around on  $SO(3)$ . This is what allows us to make a unique choice of preimage, when we are lifting a curve and following it around by demanding continuity.

Apropos lifts, a theorem.

Thm. Let  $\alpha: I \rightarrow M$  be a continuous path on  $M$  with  $\alpha(0) = x_0$ , and let  $\bar{x}_0$  be a choice of a point in  $p^{-1}(x_0)$ . Then there exists a unique continuous path  $\bar{\alpha}: I \rightarrow \bar{M}$  such that


9/23/08

(6)

$$\bar{\alpha}(0) = \bar{x}_0 \text{ and } p(\bar{\alpha}(t)) = \alpha(t), \quad t \in I.$$

The covering space is intuitively an "unrolling" of the original space; e.g.  $SU(2)$  is obtained by "unrolling"  $SO(3)$  once. The metaphor is especially clear in the case of the circle. We now examine  $\pi_1(S^1)$ . Previously we argued on intuitive grounds that  $\pi_1(S^1) = \mathbb{Z}$ , wrapping rubber bands around doorknobs, etc. Now we outline a more rigorous proof. Nakahara's discussion of this point is hard to follow.

Identify  $S^1$  with the unit circle in the complex plane:



Introduce map  $p: \mathbb{R} \rightarrow S^1: x \mapsto e^{ix}$ .  $p^{-1}$  has an infinite number of branches, e.g.  $p^{-1}(1) = \{\dots, -2\pi, 0, 2\pi, 4\pi, \dots\}$ .


$\mathbb{R}$  is a covering space of  $S^1$ . Wrap  $\mathbb{R}$  around in a helix that sits over  $S^1$  to make the projection more geometrical. Notice that  $\exists$  an open interval  $V$  around  $\frac{1}{0} \in S^1$  s.t.  $p^{-1}(V)$  is the union of an  $\infty$  number of preimages  $V_\alpha$ , disjoint. Each local part of the circle "looks like" the local part of  $\mathbb{R}$ .

9/23/08

(7)

in a neighborhood of each preimage.

Thus,  $\mathbb{R}$  is a covering space of  $S'$ , in fact it is the universal cover since  $\pi_1(\mathbb{R}) = \{e\}$  (since  $\mathbb{R}$  is contractable).



Now let  $\alpha: I \rightarrow S'$  be a loop on  $S'$  based at 1, so

$$\alpha(0) = \alpha(1) = 1.$$

Let  $\bar{\alpha}: I \rightarrow \mathbb{R}$  be the lifted loop starting at  $\bar{\alpha}(0) = 0$  in  $\mathbb{R}$ . Since  $\alpha(1) = 1$ ,  $\bar{\alpha}(1)$  must belong to  $p^{-1}(1)$ , i.e.  $\bar{\alpha}(1)$  must  $= 2n\pi$  for some  $n \in \mathbb{Z}$ . Note that  $\bar{\alpha}$  is not a loop if  $n \neq 0$ .

Now let  $\alpha, \alpha'$  be two loops on  $S'$  constructed as above with lifts  $\bar{\alpha}, \bar{\alpha}'$  both starting at 0 and ending at  $2\pi n$  and  $2\pi n'$ . Then we have that  $\alpha \sim \alpha'$  iff  $n = n'$ .

**Proof.** First suppose  $\alpha \sim \alpha'$ . Then there is a homotopy that deforms  $\alpha$  into  $\alpha'$ . Lift each curve of the deformation; this will create a homotopy deforming  $\bar{\alpha}$  into  $\bar{\alpha}'$ . But the endpoint of  $\bar{\alpha}$  or  $\bar{\alpha}'$  cannot change discontinuously, and so cannot

9/23/08

(8)


change at all. Thus  $n=n'$ . Conversely, suppose  $n=n'$ , so the lifted curves  $\bar{\alpha}, \bar{\alpha}'$  start and end at the same points. Since  $\mathbb{R}$  is simply connected, any two curves connecting the same points can be continuously deformed into one another. Do this, and use  $p$  to project the deformation onto  $S'$ , whereupon we obtain a deformation of  $\alpha$  into  $\alpha'$ . Thus  $\alpha \sim \alpha'$ .

Thus we conclude that there is a one-to-one, onto mapping between homotopy classes in  $\pi_1(S')$  and the integers  $\mathbb{Z}$ . With a little more effort, we can show that multiplication of homotopy classes corresponds to addition in  $\mathbb{Z}$ . Thus,

$$\pi_1(S') = \mathbb{Z}.$$

$\mathbb{R}$  is the universal covering space of  $S'$ ;  $S'$  has been "unrolled" an  $\infty$  number of times.

It is not necessary to unroll all the way. Consider the single edge  $ABA$  of the Möbius strip, itself a circle, and the center line  $CC$ , which is also a circle.



Define  $p: ABA \rightarrow CC$ , and  $ABA$  becomes a covering space of  $CC$ . One circle covers another (twice).

9/23/08

(9)

In another example, the 2-torus  $T^2$  is a double cover of the Klein bottle. Finding the map  $p: T^2 \rightarrow \text{Klein}$  will be left as an exercise. The 2-torus itself has a universal cover,  $p: \mathbb{R}^2 \rightarrow T^2$ . The plane is divided into parallelograms, with opposite sides identified.

It turns out that every connected space  $M$  has a covering space, although if  $M$  is simply connected then its only cover is  $M$  itself (covering once). If  $M$  is not simply connected, then it possesses one or more covers. The following is a construction of  $\bar{M}$ , the universal covering space of  $M$ .


Let  $M$  be connected and  $x_0 \in M$ . Let  $G = \pi_1(M, x_0)$ , so elements  $g \in G$  are equivalence classes of loops based at  $x_0$ .

Let  $(x, \gamma)$  be a (point, path) pair, where  $x \in M$  and  $\gamma$  is a continuous path  $:[0,1] \rightarrow M$ , with  $\gamma(0) = x_0, \gamma(1) = x$ . It is redundant to write  $(x, \gamma)$ , since  $x = \gamma(1)$ , but it is notationally convenient. Then let  $(x, \gamma) \sim (x', \gamma')$  if  $x = x'$  and  $\gamma$  homotopic to  $\gamma'$ . This is an equivalence relation. Define  $\bar{M}$  as the space of all equivalence classes,

$$\bar{M} = \{ [ (x, \gamma) ] \mid x \in M, \gamma \text{ a path } x_0 \rightarrow x \}.$$

Also, define  $p: \bar{M} \rightarrow M: [(x, y)] \mapsto x$ .

9/23/08

 $(M)$ 

$$[(x, y)] = [(x, y')] = \text{one point of } \bar{M}$$

$$[(x, y'')] = \text{a different point of } \bar{M}$$

We want to show that  $\bar{M}$  is the universal cover of  $M$ , but first it is better to get some feel for what  $\bar{M}$  looks like.

To specify a point of  $\bar{M}$  we must first choose a point  $x \in M$  and then an equivalence class of curves connecting  $x_0$  to  $x$ . The projection  $p$  just throws away the information regarding the equiv. class and returns  $x$ .


It turns out the equivalence classes of paths connecting  $x_0$  to  $x$  can be labelled by elements of  $G = \pi_1(M, x_0)$ . This is obvious in the case  $x = x_0$ , since then the paths are loops based at  $x_0$  and the equiv. classes of paths connecting  $x_0$  to  $x$  are the elements of  $G$ . But it is true when  $x \neq x_0$ , too. Since  $G$  is a discrete group, the specification of a point  $\bar{x} \in \bar{M}$  requires a point  $x \in M$  plus a choice from a discrete set (i.e.  $G$ ).

To label the homotopic equivalence classes of paths  $\gamma$  taking  $x_0 \rightarrow x$ , let  $\Gamma$  be a standard (fixed) path from  $x_0$  to  $x$  and

9/23/08

(11)

let  $\gamma$  be any path  $x_0 \rightarrow x$ :



Then associate paths  $\gamma$  ( $x_0 \rightarrow x$ ) with loops  $\alpha$  (based at  $x_0$ ) by

$$\alpha = \gamma * \Gamma^{-1}.$$


Then it's easy to see that  $\gamma \sim \gamma'$  (homotopic, as in picture above, but  $\gamma$  not  $\sim \gamma''$ ) iff  $\alpha \sim \alpha'$ . Thus, equivalence classes of paths  $\gamma$  ( $x_0 \rightarrow x$ ) can be labelled by equivalence classes of loops  $\alpha$  (based at  $x_0$ ), i.e., by elements of  $G = \pi_1(M, x_0)$ . Therefore the number of preimages of  $p^{-1}(x)$ , for any  $x \in N$ , is the number of elements in  $G$  (the order of  $G$ ). (This may be infinite.) The number of preimages of  $p^{-1}(x)$  under  $p$  is independent of  $x \in N$ ; it is "the number of times"  $M$  covers  $N$ .

But note that the labelling of points of  $p^{-1}(x)$  by elements of  $G$  depends on the choice of the standard path  $\Gamma$ . Thus there arises the question of whether this labelling is the "same" as we had for points of  $p^{-1}(x_0)$ , that is, is it continuously connected with the labelling we used at  $x_0$ ?

Here is a partial answer. It turns out we can label points of  $p^{-1}(x)$  by elements of  $G$  in a continuous manner over any simply connected region of  $\subset M$ . Let  $V$  be such a region ( $x_0$  need not be in  $V$ ), let  $x_1$  be a chosen point in  $V$ , and

9/23/08

let  $\Gamma$  be a standard path going from  $x_0$  to  $x_1$ .



Now let  $x$  be any point in  $V$  and let  $\tau_x$  be a path from  $x_1$  to  $x$ . We wish to use this to label points of  $p^{-1}(x)$ , i.e. equiv. classes of curves  $\gamma$  ( $x_0 \rightarrow x$ ). Again associate a path  $\gamma'$  ( $x_0 \rightarrow x$ ) with a loop  $\alpha$  based at  $x_0$  by

$$\alpha = \gamma * \tau_x^{-1} * \Gamma^{-1}.$$

Again,  $\alpha \sim \alpha'$  iff  $\gamma \sim \gamma'$ . This follows since  $\Gamma$  is fixed, and since  $V$  is simply connected, any 2 paths  $\tau_x, \tau'_x$  ( $x_1 \rightarrow x$ ) are homotopic. Thus, not only are points of  $p^{-1}(x)$  labelled by elements of  $G$  (as before), but the labelling is continuous as  $x$  moves around inside  $V$ . This shows that

$$p^{-1}(V) \cong V \times G,$$

which, since  $G$  is a discrete set, means that  $p^{-1}(V)$  consists of a discrete set of sets  $V_g$  ( $g \in G$ ) which are disjoint and homeomorphic to  $V$ . So  $\bar{M}$  is a covering space, by the definition.

However, the continuous labelling of points of  $p^{-1}(x)$  by elements  $g \in G$  cannot be extended to all of  $M$  (unless it is simply connected, in which case  $G$  has only one element and  $\bar{M} \cong M$ .) So, in general,  $\bar{M} \not\cong M \times G$ . In fiber bundle language, we would say that the fiber bundle is nontrivial.

9/23/08

(13)

Here are some exercises if you want to understand this construction better.

1. Show that the branches (points) of  $p^{-1}(x)$  can be labelled continuously by elements of  $G$  as  $x$  moves along a path ~~not~~ starting at  $x_0$ , as long as the path does not cross itself. What happens if the path does cross itself (what happens to the labelling)? In particular, suppose the path returns to  $x_0$ , making a loop?

2. Show that  $\tilde{M}$  is simply connected, hence the universal covering space.

One more remark. If  $M$  is a group manifold, then  $\tilde{M}$  (the universal covering group) can be given the structure of a group, and  $p$  is a homomorphism. This is the relation between  $SU(2)$  and  $SO(3)$ , and  $\mathbb{R}$  and  $S^1$ .

9/23/08

14

Now we develop notation for groups, in order to talk about  $\pi_1(M)$  for various  $M$ .

Word

Let  $G$  be a group and  $X = \{a, b, c, \dots\}$  a finite collection of elements of  $G$ . If every element  $g \in G$  can be written as products of powers of  $a, b, c, \dots$ , written in some order, including negative powers, then  $G$  is said to be finitely generated and  $X$  to be the set of generators. For example, we might have  $aba^{-1}$ ,  $a b^{-2} c b^2 a^3$ , etc. If two identical generators are adjacent, they may be combined, e.g.  $a b^2 a a^{-2} c = a b^2 a^{-1} c$ . If any generator to the zero power occurs, we drop it, since  $a^0 = e = 1$  = identity. A product of generators obeying these rules will be called reduced. If every element of  $G$  can be written uniquely as a reduced product of generators, then  $G$  is said to be freely generated.

If  $G$  is ~~not~~ finitely generated by  $X \subset G$  but not freely, then there are relations connecting the generators. For example, an Abelian group with two generators  $X = \{a, b\}$  will satisfy the relation  $ab = ba$ . This means that elements  $g \in G$  can be written as reduced products in more than one way. We handle this case by going back to the free group, and then dividing by a subgroup.

Let  $F$  be the free group constructed out of  $n$  generators  $(x_1, \dots, x_n)$ . Think of the generators as an alphabet. A word is a product of powers of letters,

$$x_{j_1}^{i_1} x_{j_2}^{i_2} \dots x_{j_s}^{i_s},$$

where  $i_k \in \mathbb{Z}$  and ~~not zero~~.  $1 \leq j_k \leq n$ . If a word is reduced according to the rules above, then it is said to be reduced. Let  $F$  = set of reduced words.  $F$  acquires the structure of a group when we define

9/23/08

(15)

the product

multiplication by concatenating, then reducing ~~words~~. The identity is the word of length zero.

Suppose  $G$  is generated by  $\{x_1, \dots, x_n\}$  but not freely. Then we can define a map

$$f: F \rightarrow G$$

which just maps reduced words in  $F$  into the same expression in  $G$ .

However, this map is not injective if  $G$  is not free, i.e., there will be more than one <sup>reduced</sup> word in  $F$  that gives rise to a given element of  $G$  (for example,  $xy$  and  $yx$  if  $G$  is Abelian). The map is, however, surjective, since we are assuming the  $\{x_1, \dots, x_n\}$  generate  $G$ . The map  $f$  is also a group homomorphism, because the rules for combining words in  $F$  ~~is~~ also work in  $G$ . Therefore  $G = F / \ker f$ , where  $\ker f$  is the normal subgroup of  $F$  that is mapped onto the identity in  $G$ . ~~Words in  $\ker f$  have the form  $gfg^{-1}$ , where it is a relation~~

$\ker f$  is specified by constraints on the generators, also called relations. Some examples will give the idea.

Let  $G$  = Abelian group, gen. by  $\{x, y\}$ . The relation is  $xyx^{-1}y^{-1} = 1$ . We write  $G = \{x, y; xyx^{-1}y^{-1}\}$  as the presentation of  $G$ . Another example,  $G = \mathbb{Z}_k$ . Here we have one generator and one relation,  $G = \{x; x^k\}$ .


More formally,  $C = \text{constraint subgroup of } G$  is defined by

$$C = \text{gen} \{ grg^{-1} \mid g \in G, r \in R \}$$

where  $R \subseteq G$  is the set of relations.  $C$  is a normal subgroup

Now some more homotopy groups. First the torus:

9/23/08




all 4 corners are same point.

Square with opposite sides identified. Obvious guesses for generators of  $\pi_1(T^2)$  are loops A, B. But clearly  $ABA^{-1}B^{-1}$  is contractible, so the group is

$$\pi_1(T^2) = \{A, B; ABA^{-1}B^{-1}\} = \mathbb{Z}^2.$$

Next Klein bottle. Square with a twisted identification.



again, all 4 corners are same pt.

This time  $ABAB^{-1}$  is contractible, and

$$\pi_1(\text{Klein bottle}) = \{A, B; ABAB^{-1}\}. = \text{non-Abelian.}$$

Now remark on relation between homotopy and homology. The relation involves the commutator subgroup of an arbitrary group, which I now explain.

Let G be any group and let C be generated by all elements of the form,

$$xyx^{-1}y^{-1}, x, y \in G.$$

$C$  called the commutator of 2 group elements.

$C$  is called the commutator subgroup.

9/23/08

Thm:  $C$  is normal. Proof: Let  $g \in G$ , then

$$\begin{aligned} gxyx^{-1}y^{-1}g^{-1} &= (gxg^{-1})(gyg^{-1})(gx^{-1}g^{-1})(gy^{-1}g^{-1}) \\ &= x'y'x'^{-1}y'^{-1} \quad \text{where } x' = gxg^{-1} \\ &\quad y' = gyg^{-1}. \end{aligned}$$

Hence  $gCg^{-1} = C$ ,  $\forall g \in G$ . Thus the quotient group  $G/C$  is defined.

Thm:  $G/C$  is Abelian. It is a kind of Abelianized version of  $G$ .

Proof: Let  $[g_1], [g_2]$  be two cosets of  $C$ . Then

$$g_1g_2 \underbrace{(g_2^{-1}g_1^{-1}g_2g_1)}_{\hookrightarrow \in C} = g_2g_1,$$

so  $[g_1g_2] = [g_2g_1] = [g_1][g_2] = [g_2][g_1]$ ,  $\frac{G}{C}$  is Abelian.

Thm: Let  $K$  be a simplicial complex, let  $G = \pi_1(K)$  and  $C$  = commutator subgp. of  $G$ . Then

~~$$H_1(K, \mathbb{Z}) \cong \frac{\pi_1(K)}{C}.$$~~

Proof omitted.

E.g. with Klein bottle,  $\pi_1(M) = \{x, y; xyx^{-1}y^{-1}\}$ .

To divide by commutator subgroup, append extra relation  $xyx^{-1}y^{-1}$ .

If  $xyx^{-1}y^{-1} = 1$  and  $xy = yx$ , then  $x(yx)y^{-1} = x(xy)y^{-1} = x^2 = 1$ .

Thus,  $H_1(\text{Klein})$  is the Abelian group generated by  $\{x, y\}$  with the relation  $x^2 = 1$ , i.e., it is  $\mathbb{Z} \times \mathbb{Z}_2$ .