

9/25/08

①

Now we examine the higher homotopy groups.



Recall that to study maps: $S^1 \rightarrow M$, we studied maps: $I \rightarrow M$ ($I = [0, 1] \subset \mathbb{R}$),
subjected to the condition that $f(0) = f(1) = \bullet$ (these are loops). This was a
matter of convenience.



Similarly, to study maps: $S^2 \rightarrow M$ it is convenient instead to look at maps

$$\alpha: I \times I \rightarrow M,$$

where $I \times I$ is the square and it is understood that the boundary of $I \times I$
is mapped to a single pt.



Note, square with all pts on bdry ∂I identified $\cong S^2$. Call such a map
a 2-loop. Then the rest of the story proceeds very much as in the
case of $\pi_1(M)$: More generally, consider maps $\alpha: \underbrace{I \times \dots \times I}_{I^n} \rightarrow M$
(These are n-loops.) $\alpha: \partial I \rightarrow x_0$.


1. $\alpha \sim \beta$, 2-loops α and β are homotopic, if \exists a smooth
interpolating map (the homotopy) that preserves the boundary
point (i.e., maps the bdry to x_0 for all values of the deform. param.)
2. $\alpha \sim \beta$ is an equivalence relation, hence classes $[\alpha]$, $[\beta]$
etc. meaningful.
3. $\alpha * \beta$ is defined by

$$(\alpha * \beta)(s_1, s_2) = \begin{cases} \alpha(2s_1, s_2) & 0 \leq \frac{1}{2} \leq s_1 \\ \beta(2s_1 - 1, s_2) & \frac{1}{2} \leq s_1 \leq 1. \end{cases}$$

②

9/25/08


Thus, pictorially:



4. $[\alpha] * [\beta]$ is defined ($*$ respects homotopy classes), and other axioms of a group are satisfied. The group (for n -loops) is denoted $\pi_n(M, x_0)$.
5. $\pi_n(M, x_0)$ is isomorphic to $\pi_n(M, x_1)$, if M is arcwise connected. Hence we just write $\pi_n(M)$ for the abstract group (the n -th homotopy group).
6. If X is of same homotopy type as Y , then $\pi_n(X) = \pi_n(Y)$.
7. $\pi_n(X \times Y) = \pi_n(X) \times \pi_n(Y)$.

There is one important property of the higher homotopy groups ($n \geq 2$) not shared by π_1 : The higher homotopy groups are Abelian.

Reason for this can be seen pictorially:



where the shaded region (and all boundaries) are mapped into x_0 .



9/25/08

③

This shows that $\alpha * \beta \sim \beta * \alpha$, hence $[\alpha] * [\beta] = [\beta] * [\alpha]$.

Another property of higher homotopy groups not shared by π_1 concerns the universal covering space. Let M be a space and \bar{M} its universal cover. Then $\pi_1(\bar{M}) = \{e\}$ (\bar{M} is simply connected), but in general $\pi_1(M)$ is not trivial (if it is trivial, then $M = \bar{M}$).

But for $n \geq 2$, $\pi_n(\bar{M}) = \pi_n(M)$. The basic idea behind this fact is that the spheres S^n are simply connected for $n \geq 2$, so it's possible to define the lift of an n -loop:



The lift of a point x on the 2-loop α is the equivalence class $[(x, \gamma)]$, where γ is a path from x_0 to x confined to α . Since the 2-loop α is simply connected, it means that the class $[(x, \gamma)]$ is independent of the choice of γ , and therefore specifies a unique point in \bar{M} .

As an application of this, note that $\mathbb{R}P^n$ is covered by S^n , so

$$\pi_k(\mathbb{R}P^n) = \pi_k(S^n). \quad \text{So what is } \pi_k(S^n)?$$

9/25/08

④

simplest case is $\pi_n(S^n)$, which concerns mappings of $S^n \rightarrow S^n$. Recall $\pi_1(S^1) = \mathbb{Z}$, where $n \in \mathbb{Z}$ is interpreted as a winding number. There is a generalization of this to higher dimensions, i.e., for a map $f: S^n \rightarrow S^n$ it is possible to say "how many times" the image of f "wraps around" S^n . This number is called the Brouwer index or degree of f . And, as in the case $n=1$, it turns out that the Brouwer degree uniquely characterizes the homotopy classes. Thus,

$$\pi_n(S^n) = \mathbb{Z} \quad (\text{all } n \geq 1).$$

What about the case $\pi_k(S^n)$ for $k < n$? Recall $\pi_1(S^n) = \{e\}$ for $n \geq 2$ (the loop is contractible on the face of S^n). Something like this also happens for $\pi_k(S^n)$ when $k < n$, that is,

$$\pi_k(S^n) = \{e\}, \quad 1 \leq k < n.$$

It turns out the case $k > n$ is also interesting. It's not easy to see this, because the highest dimensional case that is easy to visualize concerns maps $f: S^2 \rightarrow S^1$ ($k=2, n=1$), and all such maps are trivial (= contractible). (Think of mapping a sphere S^2 onto the equator). Thus,

$$\pi_2(S^1) = \{e\}.$$

But it turns out there are nontrivial (not homotopic to constant map) maps from S^3 to S^2 . The Hopf map discussed earlier in the

9/25/08

⑤

course is an example of one of these. In fact,

$$\pi_3(S^2) = \mathbb{Z}.$$

In all these results, $k > 1$, we can replace S^n by $\mathbb{R}P^n$.

See the table in Nakahara, p. 151.

Finally, what about the case $n=0$? A literal interpretation of the definition of $\pi_n(M, x_0)$ to the case $n=0$ would involve maps $f: S^0 \rightarrow M$ with one point fixed. S^0 is properly just 2 points (± 1 in \mathbb{B}), so we can say $f(1) = x_0$, $f(-1) = \text{another pt. } x_1$, say. But if M is connected, then x_1 can be continuously pulled to x_0 , and $\pi_0(M) = \{e\}$, all connected M .

Now go back to defects in CM systems, now that we know about higher homotopy groups. Make a table of various order parameter spaces (OPS's):

System	OPS = M
Liqu. ^4He x, y spin model	S^1
Nematic liquids	$\mathbb{R}P^2$
Dipole locked ^3He phase A	$SO(3) = \mathbb{R}P^3$

Recall, ^{in 3D} point defects are described by nontrivial homotopy classes $\pi_2(M)$, line defects (vortices) by $\pi_1(M)$, and textures (field configurations) with asymptotically constant values by $\pi_3(M)$.

9/25/08

Make a table of homotopy groups:

OPS	π_1	π_2	π_3
S^1	\mathbb{Z}	$\{e\}$	$\{e\}$
$\mathbb{R}P^2$	\mathbb{Z}_2	\mathbb{Z}	\mathbb{Z}
$SO(3) = \mathbb{R}P^3$	\mathbb{Z}_2	$\{e\}$	\mathbb{Z}
S^2	$\{e\}$	\mathbb{Z}	\mathbb{Z}

For example, a unit vector field $\hat{n}(\vec{r})$ (OPS = S^2) can possess point defects (characterized by an integer) and field configurations that are ~~with~~ asymptotically constant (also characterized by an integer), but no line defects (vortices).

Now start on manifold theory. Most of the spaces that occur in physics are differentiable manifolds. The idea is that a diff. manifold is a topological space with enough extra structure to talk about differentiability. That is, one can do calculus on manifolds.

Def: A differentiable manifold M is a topological space ~~with~~ plus a set $\{U_i, \varphi_i\}$, where $\{U_i\}$ is an open cover of M (each U_i is an open subset of M , and $\bigcup_i U_i = M$), and where φ_i is a map

$$\varphi_i: U_i \rightarrow \mathbb{R}^n$$

such that:

a) φ_i is a homeomorphism onto its image V_i in \mathbb{R}^n